



Australia's National
Science Agency

Tier 4 assessment for Blue-eye Trevalla (*Hyperoglyphe antarctica*) slope (data to 2020)

Prepared for SERAG MS Teams video meeting

29 November - 1 December 2021

Commercial in confidence

22 November 2021

Citation

Sporcic, M. (2021). *Tier 4 Assessment for Blue-eye Trevalla (Hyperoglyphe antarctica) slope (data to 2020)*. Technical report presented at the SERAG, MS Teams meeting 29 November – 1 December 2021. CSIRO Oceans and Atmosphere, Hobart. 16 p.

Copyright (2021)

© 2021 CSIRO. To the extent permitted by law, all rights are reserved, and no part of this publication covered by copyright may be reproduced or copied in any form or by any means except with the written permission of CSIRO.

Important disclaimer

CSIRO advises that the information contained in this publication comprises general statements based on scientific research. The reader is advised and needs to be aware that such information may be incomplete or unable to be used in any specific situation. No reliance or actions must therefore be made on that information without seeking prior expert professional, scientific and technical advice. To the extent permitted by law, CSIRO (including its employees and consultants) excludes all liability to any person for any consequences, including but not limited to all losses, damages, costs, expenses, and any other compensation, arising directly or indirectly from using this publication (in part or in whole) and any information or material contained in it.

CSIRO is committed to providing web accessible content wherever possible. If you are having difficulties with accessing this document, please contact csiroenquiries@csiro.au.

Use of this Report

The use of this Report is subject to the terms on which it was prepared by CSIRO. In particular, the Report may only be used for the following purposes:

- ❖ this Report may be copied for distribution within the Client's organisation;
- ❖ the information in this Report may be used by the entity for which it was prepared (-the Client-), or by the Client's contractors and agents, for the Client's internal business operations (but not licensing to third parties);
- ❖ extracts of the Report distributed for these purposes must clearly note that the extract is part of a larger Report prepared by CSIRO for the Client.

The Report must not be used as a means of endorsement without the prior written consent of CSIRO.

The name, trademark or logo of CSIRO must not be used without the prior written consent of CSIRO.

Contents

Acknowledgements	iii
Executive Summary	iv
1 Introduction	1
1.1 Tier 4 Harvest Control Rule	1
1.2 Tier 4 Assumptions.....	1
1.2.1 Informative CPUE.....	1
1.2.2 Consistent CPUE Through Time	1
1.2.3 Plausible Target Reference Period.....	2
1.2.4 Accurate Total Catch History	2
1.2.5 Some Implications of the Assumptions	2
2 Blue-eye Trevalla	3
2.1 Discussion.....	4
References	5
3 Appendix: Methods	6
3.1 Tier 4 Harvest Control Rule	6
3.2 The Inclusion of Discards.....	7
3.2.1 Analyses Including Discards	8
3.2.2 The Limitations of Including Discards.....	9
3.3 Selection of Reference Periods	9

Acknowledgements

Thanks goes to the CSIRO database team for their processing of the catch and effort (CPUE) and Catch Disposal Record (CDR) data as received from the Australian Fisheries Management Authority. Thanks also goes to Dr Geoff Liggins (NSW Department of Primary Industries) for providing revised NSW catch series for the assessed species. Geoff Tuck (CSIRO) and Jemery Day (CSIRO) are also thanked for reviewing this report.

Executive Summary

A Tier 4 assessment was performed for the following species:

- ❖ Blue-eye Trevalla slope (*Hyperoglyphe antarctica*)

The catch-time series used in this assessment was derived from Sporcic and Day (2021). Furthermore, as requested by SERAG in 2020, the standardized CPUE series was based on data corresponding to SESSF zones 20-50 and the Great Australian Bight (GAB) (Sporcic 2021). However, the standardized CPUE series used in the previous Tier 4 assessment was based on SESSF zones 20-50 only, i.e., excluding the GAB (Sporcic 2020).

The 2021 RBC was approximately 349.32 t, corresponding to a 122.29 t increase compared to the 2020 RBC, i.e., 227.03 t (Sporcic 2020). This 54% increase in RBC between assessments can be mostly attributed the use of the new standardized CPUE series which resulted in a higher most recent four-year average compared with the corresponding average standardized CPUE from the previous assessment. The scaling factor of approximately 54% which is applied to the target catch reflects this RBC-increase. The 2021 estimated RBC (i.e., for the 2022 fishing season) is greater than the reported catch of approximately 225.1 t in 2020 for this species.

1 Introduction

1.1 Tier 4 Harvest Control Rule

The Tier 4 harvest control rules are the default procedure applied to species which only have catches and catch per unit effort (CPUE) data available; specifically, there is no other reliable information on either current biomass levels or current exploitation rates.

Ideally, in line with the notion of being more precautionary in the absence of information, the outcome from these analyses should be more conservative than those available from higher Tier analyses; this is now explicitly implemented by imposing a 15% discount factor on the Tier 4 RBC as a precautionary measure unless there are good reasons for not imposing such a discount on particular species. The application of the discount factor will occur unless RAGs generate explicit advice that alternative equivalent precautionary measures are in place (such as spatial or temporal closures) or that there is evidence of historical stability of the stock at current catch levels (AFMA, 2009).

Tier 4 analyses require as a minimum, a time series of total catches and of standardized CPUE, along with an agreed reference period and reference points.

The current Tier 4 analysis and control rule underwent Management Strategy Evaluation (Wayte, 2009; Little et al., 2011a), which demonstrated its advantages over an earlier implementation used in 2007 and 2008. Further work has since demonstrated that if there is a limit on increases and decreases to the RBC of no more than 50 % then the notion of including a maximum RBC (at 1.25 times the target) is redundant (Little et al., 2011b).

1.2 Tier 4 Assumptions

1.2.1 Informative CPUE

There is a linear relationship between CPUE and exploitable biomass. If there is hyper-stability (CPUE remain stable while stock size changes) or hyper-depletion (CPUE decline much faster than stock size changes) then the standard Tier 4 analysis would provide biased results.

1.2.2 Consistent CPUE Through Time

The character of the estimated CPUE has not changed in significant ways through the period from the start of the reference period to the end of the most recent year. If there has been significant effort creep altering the catchability, or there have been changes to the fleet that have altered the relative efficiency of the vessels fishing, or the catchability of the species by the fleet has been altered by other changes then the comparability of recent CPUE with the target period may be compromised. Such changes would obviously reduce the responsiveness of the Tier 4 method to change and may generate completely inappropriate management advice. Included in this clause are the effects of targeting or not targeting of deep water or aggregated species. When CPUE are extremely variable through time, such that mean estimates become unreliable measures of stock status, then the Tier 4 approach cannot be validly applied.

1.2.3 Plausible Target Reference Period

The reference period provides a good estimate of the stock when at a depletion level of 48 % unfished spawning biomass. The Tier 4 method is based on CPUE and thus relates to exploitable biomass and not spawning biomass. As a minimum the reference period will refer to a period when the stock was in an acceptable, productive, and sustainable state. But there can be no guarantees that the target aimed for is really $B_{48\%}$.

1.2.4 Accurate Total Catch History

Accurate estimates are required for all catches from the stock under consideration during the accepted target period, irrespective of what method was used or whether it was retained or discarded. This assumption is especially vulnerable to being breached when large proportions of catches are discarded. While there is a procedure for adjusting the standardized CPUE for these missed catches the uncertainty over the actual number of fish killed remains.

1.2.5 Some Implications of the Assumptions

The outcomes of the Tier 4 analysis should not be regarded with the same confidence as those from Tier 1 assessments. Even though they are termed stock assessments, in actuality they are empirical considerations of catches and CPUE. Any uncertainty in the catch or CPUE time series is propagated directly through to the outputs of the analysis. For quota species the catches and reported CPUE is usually relatively well founded because of the quota catch disposal records and other compliance requirements. However, where there is a relatively high degree or variable discarding of catches this can lead to much greater levels of uncertainty.

The assessments for those species that are conducted using a Tier 4 analysis should be reviewed for their inter-annual consistency and how the fishery has been responding to the management advice derived from the Tier 4 assessments.

2 Blue-eye Trevalla




Figure 1: Blue-eye Trevalla. Top plot is the total removals with the fine line illustrating the target catch. Bottom plot represents the standardized CPUE with the upper fine line representing the target CPUE and the lower line the limit CPUE. Thickened lines represent the reference period for catches, CPUE, and the recent average CPUE. The thin black dotted line is the unmodified standardized CPUE. Discards are assumed to be

Table 1: Blue-eye Trevalla RBC calculations. C_{targ} and $\text{CPUE}_{\text{targ}}$ (CE_Target) are the targets identified in the figure above, CPUE_{Lim} is 20% of the B0 proxy (which relate to the $\text{CPUE}_{\text{targ}}$), and the most recent CPUE is the average CPUE over the last four years (CE_Recent). The RBC calculation does not account for predicted discards of predicted State catches.

Parameter	Value	Parameter	Value
Reference_Years	1997 - 2006	Scaling	0.5428
CE_Target	1.2287	Previous TAC (t)	448
CE_Limit	0.512	C_{targ}	643.497
CE_Recent	0.901	RBC	349.321
Wt_Discard	-		

Table 2: Blue-eye Trevalla data for the Tier 4 calculations. Total (t) is the sum of State, Non-Trawl and SEF2 catches. All values in Tonnes. CE is the standardized CPUE corresponding to zones 20-50 and the Great Australian Bight (Sporcic, 2021).

Year	Catch	Total	State	Non-Trawl	CE	TAC
1997	821.73	821.73	620.21	205.86	1.8588	125
1998	595.45	595.45	121.36	380.44	1.5397	630
1999	676.58	676.58	132.61	464.66	1.5036	630
2000	747.77	747.77	89.46	567.19	1.2457	630
2001	653.47	653.47	78.18	478.40	1.2633	630
2002	553.90	553.90	102.36	427.97	1.0143	630
2003	555.19	555.19	55.73	556.56	0.9243	690
2004	693.34	693.34	66.87	566.92	1.0915	621
2005	543.71	543.71	62.94	449.20	0.8243	621
2006	593.84	593.84	45.61	496.74	1.0213	560
2007	643.24	643.24	57.79	536.28	1.2025	785
2008	411.15	411.15	37.78	338.85	0.8814	560
2009	467.25	467.25	38.76	404.11	0.9696	560
2010	430.73	430.73	47.86	358.81	0.6305	428
2011	422.53	422.53	46.25	430.06	0.7252	326
2012	293.34	293.34	34.52	307.37	0.7197	388
2013	287.90	287.90	24.05	252.18	0.8868	388
2014	339.64	339.64	21.15	292.21	1.1075	335
2015	259.40	259.40	23.68	267.52	1.0532	335
2016	253.36	253.36	16.70	310.36	0.9480	410
2017	374.91	374.91	19.32	355.62	0.9381	458
2018	361.39	361.39	23.85	305.37	1.0071	462
2019	299.42	299.42	9.40	277.61	0.8724	458
2020	225.09	225.09	9.42	211.26	0.7865	448

2.1 Discussion

The catch-time series used in this assessment (Table 1) was derived from Sporcic and Day (2021). Furthermore, as requested by SERAG in 2020, the standardized CPUE series was based on data corresponding to SESSF zones 20-50 and the Great Australian Bight (GAB) (Table1; Sporcic 2021). However, the standardized CPUE series used in the previous Tier 4 assessment was based on SESSF zones 20-50 only, i.e., excluding the GAB (Sporcic 2020).

The 2021 RBC was approximately 349.32 t (Table 1), corresponding to a 122.29 t increase compared to the 2020 RBC, i.e., 227.03 t (Sporcic 2020). This 54% increase in RBC between assessments can be mostly attributed the use of the new standardized CPUE series which resulted in a higher most recent four-year average compared with the corresponding average standardized CPUE from the previous assessment. The scaling factor of approximately 54% which is applied to the target catch reflects this RBC-increase. The 2021 estimated RBC (i.e., for the 2022 fishing season) is greater than the reported catch of approximately 225.1 t in 2020 for this species.

References

- AFMA (2009). SESSF Stock Assessment Methods and TAC Setting Process Version 1.5. 8 p.
- AFMA (2017). *Southern and Eastern Scalefish and Shark Fishery Management Arrangements Booklet 2017*. Australian Fisheries Management Authority, Canberra, Australia. 92 p.
- Haddon, M. (2010). Tier 4 Analyses (data from 1986 - 2008). Pp 319 - 369 in Tuck, G.N. (ed) *Stock Assessment for the Southern and Eastern Scalefish and Shark Fishery 2009 Part 2*. Australian Fisheries Management Authority and CSIRO Marine and Atmospheric Research, Hobart. 428 p.
- Haddon, M. (2014). Catch Rate Standardizations for Selected Species from the SESSF (data 1986-2012) pp 57-275 in Tuck, G.N. (ed) (2014). *Stock Assessment for the Southern and Eastern Scalefish and Shark Fishery 2013. Part 2*. Australian Fisheries Management Authority and CSIRO Marine and Atmospheric Research, Hobart. 487 p.
- Haddon, M., Sporcic, M. (2017). *Tier 4 Assessments for selected SESSF Species (data to 2016)*. CSIRO Oceans and Atmosphere, Hobart. 52 p.
- Little, R., Tuck, G.N., Haddon, M., Day, J., Klaer, N., Smith, A.D.M., Thomson, R., Wayte, S. (2009). Developing CPUE targets for the Tier 4 harvest strategy of the SESSF. Pp 233-254 in Tuck, G.N. (ed) *Stock Assessment for the Southern and Eastern Scalefish and Shark Fishery: 2008 Part 2*. Australian Fisheries Management Authority and CSIRO Marine and Atmospheric Research, Hobart. 331 p.
- Little, L.R., Wayte, S.E., Tuck, G.N., Smith, A.D.M., Klaer, N., Haddon, M., Punt, A.E., Thomson, R., Day, J., Fuller, M. (2011a). Development and evaluation of a cpue-based harvest control rule for the southern and eastern scalefish and shark fishery of Australia. *ICES Journal of Marine Science* 68(8): 1699-1705.
- Little, L.R., Wayte, S.E., Tuck, G.N. (2011b). The effects of Cmax on the Tier 4 Harvest Control Rule. Pp 3-9 in Tuck, G.N. (ed) *Stock Assessment for the Southern and Eastern Scale-fish and Shark Fishery: 2010 Part 2*. Australian Fisheries Management Authority and CSIRO Marine and Atmospheric Research, Hobart. 419 p.
- R Core Team (2021). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
- Sporcic, M. (2020). *Tier 4 Assessments for selected SESSF species (data to 2019)*. Technical report presented to the SERAG, Dec 9-10, 2020. CSIRO Oceans and Atmosphere, Hobart. 37 p.
- Sporcic, M. (2021). *Update Part 2: Statistical CPUE (catch-per-hook) Standardizations for Blue-eye Trevalla (Auto-line and Drop-line) in the SESSF (data to 2020)*. CSIRO, Hobart. Technical report presented at SERAG meeting 29 November to 1 December 2021. 26 p.
- Sporcic, M., Day J. (2021). *Draft catch history time series for selected Tier 4 SESSF species (data to 2020)*. Prepared for SERAG meeting 28-29 September 2021. 19 p.
- Wayte, S.E. (ed.) (2009). *Evaluation of new harvest strategies for SESSF species*. CSIRO Marine and Atmospheric Research, Hobart and Australian Fisheries Management Authority, Canberra. 137 p.

3 Appendix: Methods

3.1 Tier 4 Harvest Control Rule

The data required are time series of catches and standardized CPUE. The analyses have been conducted on total catches across the entire SESSF (including State catches, SEF2 landing records, and any discards). For some species, where there is only a single stock and a single primary fishing method, analyses are presented using standardized CPUE data (e.g., Haddon, 2014). For other species, there may be multiple stocks or areas or multiple methods and selecting which time series of CPUE to use in the analyses is not always straightforward. In those cases, the standardized CPUE time series for the method now accounting for the majority of current catch was used.

All 2010 data relating to catches and discards, from both State waters and SEF2 data sets, were provided by AFMA, with initial processing by N. Klaer and J. Upston of CSIRO. All CPUE data were derived from the standard commercial catch and effort database processed by the data services Team at CSIRO Hobart.

Standard analyses were set up in the statistical software, R Core Team (2021), which provided the tables and graphs required for the Tier 4 assessments. The data and results for each analysis are presented for transparency. The Tier 4 harvest control rule formulation essentially uses a ratio of current CPUE with respect to the selected limit and target reference points to calculate a scaling factor for the current year. This scaling factor is applied to the target catch to generate an RBC. To generate a TAC, known discards and State catches are first removed and then, if applicable, the 15% discount is applied. The TAC calculations are conducted by AFMA. This report focusses on providing the estimates of the Recommended Biological Catches.

$$\text{Scaling Factor} = SF_t = \max\left(0, \frac{\overline{CPUE} - CPUE_{lim}}{CPUE_{targ} - CPUE_{lim}}\right)$$
$$RBC = C_{targ} \times SF_t$$

If new data becomes available, for example, more State data has become available this year, or other large changes occur in the CPUE then the RBC could undergo large changes. Such changes are constrained by the following limits:

$$RBC_y = 1.5RBC_{y-1} \quad RBC_y > 1.5RBC_{y-1}$$
$$RBC_y = 0.5RBC_{y-1} \quad RBC_y < 0.5RBC_{y-1}$$

where

1. RBC_y is the RBC in year y ,
2. $CPUE_{targ}$ is the target CPUE for the species,
3. $CPUE_{lim}$ is the limit CPUE for the species = $0.4 * CPUE_{targ}$,
4. \overline{CPUE} is the average CPUE over the past m years; m tends to be the most recent four years,
5. C_{targ} is a catch target derived from a period of historical catch that has been identified as a desirable target in terms of CPUE, catches and status of the fishery, e.g. 1986 – 1995. This is an average of the total removals for the selected reference period, including any discards.

$$C_{targ} = \frac{\sum_{y=yr1} L_y}{(yr2 - yr1 + 1)}$$

where L_y represents the landings in year y .

$$CPUE_{\text{targ}} = \frac{\sum_{y=yr1}^{yr2} CPUE_y}{(yr2 - yr1 + 1)}$$

where $CPUE_y$ is the CPUE in year y , $yr2$ and $yr1$ represent the last and the first years in the reference period respectively.

Percent discards are estimated from ISMP observations from 1998 to the current year. Discards for earlier years, prior to ISMP sampling, are generally estimated by taking the overall average percent discard from 1998 to the 2006 and applying that discard rate to the reported landings for the earlier years. The year 2006 was selected as the final year as discarding practices altered at about that time following the structural adjustment and the introduction of the Harvest Strategy Policy. For Eastern Gemfish the average discard rate was determined for 1998-2002 to allow for the non-target nature of the fishery following 2002. The calculation of the earlier discards is done so that the total catches can be estimated even though only the landed catches are available. To calculate the discards for a given year we used:

$$D_y = \frac{C_y \bar{D}_{98-06}}{(1 - \bar{D}_{98-06})}$$

Discard proportions for the projected year for which the RBC is being calculated are taken as a weighted mean of the previous four years:

$$D_{\text{CUR}} = (1.0 D_{y-1} + 0.5 D_{y-2} + 0.25 D_{y-3} + 0.125 D_{y-4})/1.875$$

where D_{CUR} is the estimated discard rate for the coming year y , D_{y-1} is the discards rate in year $y-1$. The discard rate in year y is the ratio of discards to the sum of landed catches plus those discards (this can vary between 0 – 100 %):

$$D_y = \frac{\text{Discard}_y}{(\text{Catches}_y + \text{Discard}_y)}$$

For each species, reference years were selected by the RAGs to generate estimates of target catches and target CPUE. In addition, a decision was required as to whether the fishery could be considered as fully developed or otherwise. Where a fishery was not considered to be fully developed the target CPUE, $CPUE_{\text{targ}}$, was divided by two as a proxy for expected changes to CPUE as the fishery develops and the resource stock size declines towards the target of 48% unfished biomass.

Plots are given of the total removals illustrating the target catch level. In addition, the standardized CPUE are illustrated with the target CPUE and the limit CPUE. Finally, where the data are available, plots are given of the Total removals contrasted with State removals, and of discards and non-trawl catches.

3.2 The Inclusion of Discards

Some species, especially redfish (*Centroberyx affinis*) and inshore Ocean Perch (*Helicolenus percoides*), have experienced high levels of discarding but the reported CPUE relate only to the estimated landed weights. In those species where discarding makes up a significant proportion of the catch (in some years more redfish were discarded than landed and more inshore ocean perch tend to be discarded than landed) it is reasonable to ask how the discards would have affected CPUE. This is an important question because standardized commercial CPUE are used in Australian stock assessments as an index of relative abundance (e.g., Haddon, 2014); if ignoring discards leads to a consistent bias this could affect the outcome of the assessments and thus, the assessments should become aware of the effects of discards.

CPUE are used in assessments as an index of relative abundance through time and it is the trends exhibited by the CPUE that are important rather than their absolute values. If the discard levels are relatively

constant through time and evenly distributed amongst the fleet, then their inclusion would not be expected to influence the trends in CPUE except to add noise. In all cases the discard rates are estimates based on sub-sampling the fleet of vessels. That the estimates are uncertain can be seen simply by considering the summary data tables in this document; where discards rates are not low they are very variable between years. Redfish provide an extreme where in 1998 the estimate was 2324 t, which was nearly 56 % of the total catch, while in 1999 discards estimated at only 69 t, making up on about 5 % of the total catch. So in those cases where discard levels are low, adding discards to the estimation of CPUE is not expected to alter outcomes.

For those species, such as redfish and ocean perch, where discard rates are much higher it was decided to include those estimated catches to determine their effect on the outcome of the Tier 4 analyses. In 2010 it was concluded that while the inclusion of discards contributed a great deal of noise to the analyses, for those species where discarding made up significant proportions of the overall catch the discard augmented CPUE should be examined each year as a sensitivity analysis to contrast with the outcome from the un-augmented CPUE (Haddon, 2010).

3.2.1 Analyses Including Discards

Discard rates cannot simply be added to known catches on the way to calculating CPUE. The standardized CPUE are estimated from individual catch and effort records but the estimates of discards are summary estimates for each fishery. While a method for incrementing the standardized CPUE has been developed it should be noted that this ignores all complications relating to unknown aspects of discarding behaviour (e.g., Is the discard rate constant across all catch sizes, across all vessels, across all areas?). This means that including discard catches into the annual CPUE estimates introduces an unknown amount of uncertainty into the analysis. It should also be noted that the discard estimates are highly variable from year to year and derive from relatively small samples of all trips contributing to catches.

The method developed was to find the multiplier needed to adjust ratio mean CPUE and apply that to the standardized CPUE (Haddon, 2010). The ratio mean CPUE require the annual sum of catches for the fishery along with the sum of effort and ratio means calculated for each year. The discard estimates from the fishery can be added to the catch totals and new ratio means calculated and compared. The multiplier needed to make the same changes to the ratio mean CPUE can then be developed and applied to the standardized CPUE.

The ratio mean is simply the sum of all catches divided by the sum of effort

$$\hat{I}_{R,t} = \frac{\sum C_t}{\sum E_t}$$

where $\hat{I}_{R,t}$ is the ratio mean CPUE for year t , $\sum C_t$ is the sum of landed catches in year t , and $\sum E_t$ is the sum of effort (as hours trawled) in year t . If $\sum D_t$ is the sum of discards in year t then the discard incremented ratio mean CPUE would be:

$$\hat{I}_{D,t} = \frac{\sum C_t + \sum D_t}{\sum E_t}$$

The same values of $\hat{I}_{D,t}$ can also be obtained using the following multiplier:

$$\hat{I}_{D,t} = [(\sum D_t / \sum C_t) + 1] \times I_t$$

where I_t is the CPUE estimate to be modified by the inclusion of discards. If this is the ratio mean, then the augmented CPUE would be identical to the first equation dealing with $\sum D_t$. In practice, the CPUE used with the multiplier are the standardized CPUE (e.g. Haddon, 2014; Sporcic, 2021).

3.2.2 The Limitations of Including Discards

The discard rates are estimated as the proportion of the total catch (= landed catch plus discards), which means that discard proportions greater than 0.5 imply that more fish are discarded than landed. To calculate the discarded catches from a discard rate and the landed catches we use:

$$D_t = \left(\frac{C_t}{1 - P_t} \right) - C_t$$

where D_t is the discarded catches in year t , C_t is the total landed catches in year t , and P_t is the proportion of discards in year t . Because the divisor is $1 - P_t$ as P_t tends to 1.0 the divisor becomes very small and hence acts as a multiplier on total landed catch C_t . The effect of this is that when P_t is estimated to be above 0.5 the multiplying effect in the calculation of discards becomes grossly exaggerated (Figure 8).

It is recommended that once discard proportions are estimated to be above 0.5 or 0.6 then attention needs to be paid to whether or not the inclusion of discards into the CPUE and the calculation of the RBC can be considered valid. In such cases, for example Inshore Ocean Perch, the Tier 4 analysis may need to be rejected and some alternative adopted.




Figure A1: The influence of the proportion discarded on estimates of discarded catches. As the proportion of discards approaches 1.0 the multiplying effect in the estimation of discard amounts becomes greatly amplified.

3.3 Selection of Reference Periods

The Tier 4 requires a reference period to be selected to establish target and limit levels of CPUE and associated target levels of catch that are deemed by the RAG to act as a proxy for the desired state for the fishery. These act as a proxy for the Harvest Strategy Policy reference points of 48% and 20% unfished spawning biomass. The original Tier 4 rule that used a linear regression of the last four year's CPUE to determine whether catches increase, or decrease was not able to rebuild a resource towards a desired target level and the current approach was developed to be able to manage a fishery towards a target and away from a limit.


The essence of the Tier 4 control rule is that it sets a RAG agreed target CPUE, which has an associated target catch. An estimate of current CPUE (usually the average of the last four years) is compared with the target and a multiplier is estimated which is to be applied to the target catch to generate the recommended biological catch.

To select a reference period requires a time series of comparable CPUE. For this reason the use of standardized CPUE should be an improvement over using, for example, the observed arithmetic or geometric mean CPUE. CPUE data is available in the SESSF for all targeted species from 1986 - 2011, although it needs to be noted that the character of the fishery has changed markedly during that period. Little et al. (2009) provide a discussion on how reference periods might be selected. They proposed a default 10-year period of 1986 – 1995, stating: “We have assumed that the average CPUE from 1986 to 1995 corresponds to that which would be attained if the stock were at the level that provides the maximum economic yield, B_{MEY} . The limit CPUE is 40% of this CPUE.” (Little et al., 2009, p 234).

For each species, reference years were selected by the RAGs to generate estimates of target catches and target CPUE. In addition, a decision was required as to whether the fishery could be considered as fully developed or otherwise during the reference period or not. Where a fishery was not considered to be fully developed the target CPUE, $CPUE_{targ}$, was divided by two as a proxy for expected changes to CPUE as the fishery develops and the resource stock size declines towards the assumed proxy target for 48% unfished biomass.

Little et al. (2009) proposed three rules used to estimate the CPUE target:

1. The CPUE target for stocks fully exploited at or prior to 1986 is based on the average CPUE from 1986-1995.
2. Where fishing exploitation up to 1986 is thought to be minimal, the CPUE determined in Step 1 is halved (to provide a CPUE proxy for B_{MEY}).
3. Where fishing exploitation after 1986 is low, the first year in which catches are above 100 t signifies the start of the 10-year period for which CPUE targeted is calculated.



As Australia's national science agency and innovation catalyst, CSIRO is solving the greatest challenges through innovative science and technology.

CSIRO. Unlocking a better future for everyone.

Contact us

1300 363 400
+61 3 9545 2176
csiroenquiries@csiro.au
www.csiro.au

For further information

CSIRO
Oceans and Atmosphere
Miriana Sporcic
+61 3 6232 2222
Miriana.Sporcic@csiro.au